



Stanford
Center for International
Development

Working Paper No. 405

Estimating Price Elasticities with Non-Linear Errors in

Variables by

Aprajit Mahajan

December 2009



Stanford University

579 Serra Mall @ Galvez, Landau Economics Building, Room 153

Stanford, CA 94305-6015

ESTIMATING PRICE ELASTICITIES WITH NON-LINEAR ERRORS IN VARIABLES*

Aprajit Mahajan[†]

December 2009

Abstract

This paper estimates a price elasticity using a flexible demand specification on survey data where prices are observed with error are correlated with household characteristics. The demand function is modeled as being a polynomial (and more generally also including trigonometric terms) in the unobserved true prices and the form of the dependency between the observed (mismeasured) prices and household characteristics is modeled parametrically. I identify and estimate the model by adapting the approach of Hausman, Newey, Ichimura, and Powell (1991) and Schennach (2004). The flexible specifications allow us to observe that price elasticities vary across the price distribution, something missed in previous work using linear demand specifications.

Keywords: Price elasticities; Non-linear errors.

JEL Classification No.: C2; O0.

*I would like to thank Han Hong, Bo Honor´ and Elie Tamer for their tireless encouragement and advice. I have also benefited enormously from discussions with Angus Deaton, Ernst Schaumburg and members of the Princeton Microeconometrics Reading Group.

[†] Department of Economics, Stanford University. Mailing Address: 579 Serra Mall, Stanford, CA 94305-6072. Phone: (650) 723-3864. Fax: (650) 725-5702. Email: amahajan@stanford.edu

Estimating Price Elasticities with Non-Linear Errors in Variables*

Aprajit Mahajan[†]
Department of Economics,
Stanford University

January 2008

Abstract

This paper estimates a price elasticity using a flexible demand specification on survey data where prices are observed with error are correlated with household characteristics. The demand function is modelled as being a polynomial (and more generally also including trigonometric terms) in the unobserved true prices and the form of the dependency between the observed (mismeasured) prices and household characteristics is modelled parametrically. I identify and estimate the model by adapting the approach of [Hausman, Newey, Ichimura, and Powell \(1991\)](#) and [Schennach \(2004\)](#). The flexible specifications allow us to observe that price elasticities vary across the price distribution, something missed in previous work using linear demand specifications.

JEL Codes: C2, O0

1 Introduction

This paper contributes to the literature on the estimation of demand responses using spatial variation in prices derived from household survey data. The measure of prices obtained from such data is typically noisy and obtained in a manner so that the measurement error in prices is usually correlated with measurement error in quantities purchased. The object of interest in these exercises and in this paper is a price elasticity of demand to be estimated flexibly from a single cross section of household data when prices are measured with error. [Deaton \(1997\)](#) estimates elasticities accounting for measurement error by imposing a linear structure on the outcome equation, while [Deaton and Ng \(1998\)](#) estimate elasticities completely non-parametrically but do not deal with the measurement error problem.

This paper serves as a midway point between the previous papers by estimating the elasticities using flexible (although still parametric) functional forms while also accounting for the mismeasured

*I would like to thank Han Hong, Bo Honoré and Elie Tamer for their tireless encouragement and advice. I have also benefited enormously from discussions with Angus Deaton, Ernst Schaumburg and members of the Princeton Microeconometrics Reading Group.

[†]Mailing Address: 579 Serra Mall, Stanford, CA 94305-6072. Phone: (650) 723-3864. Fax: (650) 725-5702. Email: amahajan@stanford.edu

prices. The approach is an extrapolation of the ideas contained in Hausman, Newey, Ichimura, and Powell (1991) (henceforth HNIP) and the extensions in Schennach (2004) but incorporating the clustered feature of the data and the non-classical nature of the measurement error.

Dividing expenditure by quantity provides a measure of price which is referred to in the literature as the unit value and is available on a household basis for those households that make positive purchases. The basic idea is to use these unit values to estimate demand responses. However, unit values are not prices since they reflect in part at least the quality choices made by households (so that for instance richer households report higher unit values for the same aggregate commodity than poorer households) and also measurement errors in expenditures and/or quantities. Deaton (1987,1988) develops a framework for dealing with both problems in which the regression function for demand is modeled linearly. His results suggest that the quality issue is of smaller magnitude than the measurement error problem. Deaton and Ng (1998) calculate the price elasticity non-parametrically using the method of Hardle and Stoker (1989) but do not control for the measurement error problem.

This paper allows for a more general specification for the regression function for quantity while accounting for measurement error in a manner essentially analogous to HNIP and Schennach (2004) but at the cost of ignoring the quality issue. I model the regression function of interest as a polynomial in the (unobserved) price as well as a polynomial in expenditure to account for possible non linearities in income effects. In addition I also experiment with “Fourier” flexible functional forms for the demand function for whose estimation I need a suitable extension to the HNIP approach.

The paper is organized as follows: Section 2 describes the data at hand outlines the various demand specifications, Section 3 discusses the identification of these specifications. Section 4 uses the identification results from the previous section to outline an estimation strategy and derives the large sample properties of the proposed estimators. I then discuss the performance of the estimators in the context of monte carlo experiments and implement them on the data set at hand in Section 6 and the conclusions follow.

2 Data and Model

Before defining the model, it is instructive to describe the data since its key features will motivate the model. I use data from India’s National Sample Survey (50th Round, 1993-4) for rural Maharashtra which has information on 4440 households from 445 villages. Households are sampled using a two stage stratified sample design. In the first stage C clusters (villages) are chosen, and in the second, H households are sampled from the selected clusters. The data records household characteristics (expenditure and age-sex-caste demographics which I denote by x_{ch}) as well as total expenditure on wheat and total quantity purchased (q_{ch}) by each household. The commodity of interest is wheat,¹ a diet staple in the region, which is purchased by most households (about 82% in the survey) and

¹The figures are based on the aggregate commodity which includes market wheat, suji and bread.

its average budget share in total expenditure is about 3%.

The sampling design of the survey provides us with observations on several households within one cluster. I assume that the price of wheat is fixed within a cluster but varies exogenously across clusters so that if prices were observed I could identify the demand parameters - which are assumed constant within and between clusters - from the between cluster variation in prices. Here, I am implicitly assuming that each village constitutes a different market and that prices vary exogenously across markets and that this variation is independent of other village characteristics that influence household demand. The exogenous variation in prices can further be based on the geographic dispersion of the sampled clusters across space coupled with high transportation costs. These assumptions, while extremely stringent, are common to all exercises in estimating demand responses from survey data where prices are derived in the fashion described above and their plausibility needs to be weighed against the advantage of having a rich data set with substantial price variation, usually hard to come by in a time-series context in developing countries (see [Deaton \(1987\)](#), [Deaton \(1988\)](#)).

In addition, I must also assume that within clusters, household quantities are a small enough fraction of aggregate demand so that I can ignore simultaneity between prices and quantities in the regression specification (note that this disaggregation argument is not unproblematic, see [Kennan \(1989\)](#)) Finally, I abstract from further complex survey design issues and assume that across clusters the observations are independent and identically distributed.

The model is specified by two equations for each household – one for quantities and one for unit values (the latter is referred to as the measurement error equation). Following [Deaton \(1988\)](#), I parameterize the error equation as

$$u_{ch} = z_c + x'_{ch}\gamma_2 + \eta_{ch} \tag{1}$$

where u_{ch} is the log of the unit value for household h in cluster c , z_c is the (log of the) unobserved cluster price, x_{ch} is a vector of household characteristics including expenditure which is allowed to enter non-linearly, and η_{ch} is the household specific measurement error. The logarithmic transformation is adopted for convenience and, if $\gamma_2 = 0$, could be interpreted as implying that the error term η_{ch} is expressed (approximately) as a fraction deviation from the true price. I do, however, in subsequent work plan to experiment with untransformed unit values so that the errors will be treated as being in levels.

There is no theoretical justification for this particular parameterization and it differs from the typical (also usually atheoretical) statement of classical measurement error because of the presence of the household characteristics. Their linearity in (1) is purely pragmatic, but clearly their presence itself is important because I need to account for household heterogeneity in some fashion (since for instance richer households on average report higher unit values).

From (1) I define the “purged” unit value v_{ch} as

$$\begin{aligned} v_{ch} &\equiv u_{ch} - x'_{ch}\gamma_2 \\ &= z_c + \eta_{ch} \end{aligned} \tag{2}$$

which states that the residual from the best linear prediction of the unit value u_{ch} given household characteristics x_{ch} can be expressed as the sum of two random variables – a village specific term (z_c) which I interpret as the true (unobserved) price and a household specific term (η_{ch}) which is interpreted as measurement error.

I specify three possible demand equations. The first (D1) expresses quantities purchased as a polynomial function of the unobserved price

$$\ln q_{ch} = \sum_{j=0}^K \beta_j (z_c)^j + x'_{ch}\gamma_1 + f_c + \varepsilon_{ch} \tag{D1}$$

q_{ch} is the quantity demanded by household h in cluster c , f_c is a cluster level effect independent of both the prices and the demographics x_{ch} . K is the order of the polynomial in the unobserved price variable,² and for $K > 1$ the model is not linear in the measurement error η_{ch} so that standard instrumental variable approaches are no longer feasible. The independence of the cluster level effect and prices is quite strong but is necessary for identification. A possible scenario is that prices vary across villages solely due to transportation costs and that these variations are independent of any other village level characteristics that affect household demand.

The second specification (D2) follows [Deaton and Muellbauer \(1980\)](#)

$$w_{ch} = \sum_{j=0}^K \beta_j (z_c)^j + x'_{ch}\gamma_1 + f_c + \varepsilon_{ch} \tag{D2}$$

where w_{ch} is the budget share of wheat for the household.

The final specification (D3) follows the suggestion of [Gallant \(1981\)](#) and [Eubank and Speckman \(1990\)](#) and the regression function is modeled using a Fourier expansion by adding trigonometric functions to the quadratic terms. The simple Fourier ‘flexible’ form estimated in this paper is

$$w_{ch} = \sum_{j=0}^K \beta_j (z_c)^j + x'_{ch}\gamma_1 + \sum_{m=1}^M \beta_{K+m} \sin(mz_c) + \beta'_{K+m} \cos(mz_c) + f_c + \varepsilon_{ch} \tag{D3}$$

The justification for this specification arises both from the observation that exclusively polynomial expansions are often sensitive to outliers and the general arguments that a combination of polynomial and trigonometric terms often has desirable approximation properties (see for instance [Gallant \(1981\)](#) for an exposition on the advantages of such a specification).

²In principle, one could formulate the problem still more flexibly by letting K in all the specifications depend upon the sample size. This generalization and its implications for the empirical work is left for future research.

In all three specifications the error term ε_{ch} is independent of the random vector $\{z_c, x_{ch}\}$ and is independent across households. However, it is allowed to be correlated with η_{ch} for a given household, and this is most likely the case since measurement errors in quantities translate quite directly into measurement errors in unit values as the latter is obtained by dividing expenditure by the former. We assume that $(\varepsilon_{ch}, \eta_{ch})$ have mean zero and are independent and identically distributed across households and that all required higher moments exist. These assumptions are all stated formally below.

All three demand specifications attempt to account for the effect of household characteristics on unit values. In all of them the difference between unit value and its expectation conditional on household variables x is assumed to be representable as a sum of two independent random variables

$$v_h - \mathbb{E}(v_h|x_h) = z + \eta_h$$

In principle one could obtain $E(v|x)$ non-parametrically if the dimension of x is not too large. For the data we use, graphical displays suggest that this conditional expectation is approximately linear when x is (the log of) expenditure and household size, so we may not lose too much by directly modeling the conditional expectation as linear in x .

3 Identification

Throughout, we assume (for expositional ease) that the number of households in each village is fixed and that the econometrician has available an independent and identically distributed sample from the distribution of the random vector $\{q_h, u_h, x_h\}_{h=1}^H$ where h indexes households. We will also discuss identification assuming that the values of the parameters (γ_1, γ_2) are known so that the “purged” unit value v_{ch} is identified. This is without loss of generality, since under the assumptions stated now these objects are consistently estimable by a variety of standard techniques. For instance, under Assumption (1), the parameter vector (γ_1, γ_2) is identified

ASSUMPTION 1 *The error terms $(\varepsilon_{ch}, \eta_{ch})$ have zero mean and are uncorrelated with the household demographics $x_{ch'}$ for all households $h' \in \{1, \dots, H\}$ and the matrix $\sum_{h=1}^H \mathbb{E}\{\ddot{x}_{ch}\ddot{x}'_{ch}\}$ has full rank.³*

3.1 Identification of Specification (D1) and Specification (D2)

The basic idea for identification in the first two specifications is straightforward and we outline the basic idea before stating the result. Note that if prices were observed for each village (so that the econometrician observed an i.i.d. sample from the random vector $\{\{q_h, u_h, x_h\}_{h=1}^H, z\}$) and were independent of the village effect, the parameter vector β in each of (D1) and (D2) would be

³The double dots above a variable \ddot{x}_{ch} denote that each element in it is expressed as a deviation from the corresponding cluster mean.

identified (for instance as the probability limit of the pooled OLS estimates) since

$$\beta = \mathbb{E}(\tilde{z}_{cK}\tilde{z}'_{cK})^{-1} \mathbb{E}\left(\frac{1}{H} \sum_{h=1}^H \tilde{z}_{cK}y_{ch}\right) \quad (3)$$

where $\tilde{z}_K = (1, z, z^2, \dots, z^K)'$ where $y_h = \ln q_h - x'_h\gamma_1$ for the first specification and $y_h = w_h - x'_h\gamma_1$ for the second specification. In the case where z is not observed, β would still be identified if we could identify the outer product matrix $\mathbb{E}(\tilde{z}_K\tilde{z}'_K)$ and the vector $\mathbb{E}\left(\frac{1}{H} \sum_{h=1}^H \tilde{z}_K y_h\right)$. The basic idea behind identification is that these objects are identified as long as enough structure is placed on the error terms in the measurement error and demand specifications. This is the strategy first outlined in Hausman, Newey, Ichimura, and Powell (1991) and we modify the argument to account for the clustered nature of the data and the non-classical nature of the measurement error.

3.1.1 Identification of $\mathbb{E}(\tilde{z}_K\tilde{z}'_K)$

To illustrate, as long as the measurement error term η_h has zero expectation, $\mathbb{E}(v_{ch}) = \mathbb{E}(z_c)$ so that the mean of the price vector is identified. Next, as long as the error terms are independent across households and independent of prices, the second moment $\mathbb{E}(z_c^2)$ will be identified since for $h \neq h'$ $\mathbb{E}(v_{ch}v_{ch'}) = \mathbb{E}(z_c^2)$. In fact, as we show below, for $j < H$, the j^{th} uncentered moment of z is equal to $\mathbb{E}(v_{ch_1}\dots v_{ch_j})$ for $h_s \neq h_k$ for all $(s, k) \in \{1, \dots, j\}$. We first record this result.

ASSUMPTION 2 $\mathbb{E}(\eta_{ch}) = 0$ for all h and $(\eta_{ch}, \eta_{ch'})$ are independent of each other for all $h \neq h'$ and η_{ch} is independent of z_c

Lemma 1 Suppose that $\mathbb{E}(z_c^j)$ exists and that $j \leq H$ and that γ_2 is known. Then, under Assumption (2) $\mathbb{E}(z_c^j)$ is identified. In particular, $\mathbb{E}(v_{ch_1}\dots v_{ch_j}) = \mathbb{E}(z_c^j)$. In fact, we can consider all $\binom{H}{j}$ distinct subsets of size j from the set $\{v_1, \dots, v_H\}$ so that

$$\mathbb{E}(z_c^j) = \mathbb{E}\left[\frac{1}{\binom{H}{j}} \sum_{h_1=1}^{H-j+1} \sum_{h_2=h_1+1}^{H-j+2} \dots \sum_{h_j=h_{j-1}}^H v_{ch_1}v_{ch_2}\dots v_{ch_j}\right]$$

Since the data set for this paper typically contains about 10 households per village and the maximum order of polynomials considered is quintic, the condition $j \leq H$ is satisfied for the empirical application and so identification for second moment matrix $\mathbb{E}(\tilde{z}^K\tilde{z}^{K'})$ for the application will follow from Lemma 1.

If, however, the computation of the second moment matrix involves the calculation of $\zeta_j \equiv \mathbb{E}(z^j)$ for $j > H$, then we need to strengthen Assumption 2 and require that the higher moments of the measurement error η_{ch} exist. The ζ_j are then identified recursively using the formulae below for

$q = 0, 1, 2..$

$$\begin{aligned}\mathbb{E}\left(v_1^{q+1}v_2v_3\dots v_H\right) &= \sum_{l=0}^q \binom{q}{l} \zeta_{H+l} \mathbb{E}\left(\eta^{q-l}\right) + \sum_{l=0}^q \binom{q}{l} \mathbb{E}\left(\eta^{q-l+1}\right) \zeta_{H+l-1} \\ E\left(v_1^{q+1}\right) &= \sum_{l=0}^{q+1} \binom{q+1}{l} \zeta_l \mathbb{E}\left(\eta^{q+1-l}\right)\end{aligned}\tag{4}$$

ASSUMPTION 3 $\mathbb{E}(\eta_{ch}) = 0$ for all h and all higher moments of the error $\mathbb{E}(\eta_{ch}^l)$ and of the prices $\zeta_j \equiv \mathbb{E}(z^j)$ in the formulae above exist. In addition, $(\eta_{ch}, \eta_{ch'})$ are independent of each other for all $h \neq h'$ and η_{ch} is independent of z_c .

Under the assumption above we can identify $\mathbb{E}(z^j)$ for arbitrary $j \in \mathbb{N}$ and $j > H$ recursively. At each step q , the second equation above uses the previous information to identify $\mathbb{E}(\eta^{q+1})$ and this moment in turn is used in the first equation to identify the moment ζ_{H+q} . There are potentially several different ways to identify the ζ_j and we make no claims for the optimality of the procedure outlined above. An alternative strategy that places somewhat fewer assumptions on the higher moments of the measurement error is outlined in the appendix.

3.1.2 Identification of $\mathbb{E}(\tilde{z}_K y)$

Identification of the inner product $\mathbb{E}(z^s y)$ also follows using the same arguments as in the previous subsection. To illustrate, consider the directly identified object $\mathbb{E}(v_{h_1} y_{h_2})$ for $h_1 \neq h_2$. Then, as long as the error terms for household h_1 are independent of the error terms for the other households and are also independent of the cluster level variables (f_c, z_c) ⁴, then simple calculations show that $\mathbb{E}(v_{ch_1} y_{ch_2}) = \mathbb{E}(z_c y_{ch_2})$ for households $h_1 \neq h_2$. This result is recorded here for future reference.

ASSUMPTION 4 Suppose that the error terms $(\varepsilon_{ch}, \eta_{ch})$ have mean zero, are independent across households and are independent of (z_c, f_c) .

Lemma 2 Under Assumption (4) and for $j < H$, $\xi_j \equiv \mathbb{E}(z^j y_{ch})$ is identified and is equal to $\mathbb{E}(v_{ch_1} \dots v_{ch_j} y_{ch_{j+1}})$ where $h_s \neq h_k$ for $(s, k) \in \{1, \dots, j+1\}$

For the data set at hand, this is an adequate identification strategy since the highest cross moment needed is $\mathbb{E}(y_{ch} z_c^5)$, and as mentioned previously, there are about ten households per cluster.

3.1.3 Identification of β

If the prices are independent of the village effect in the demand equation, the parameter vector β is equal to the expression (3) each of whose two component terms are identified by the arguments

⁴In fact, the independence between the village level effect f_c and ε_{ch} is not necessary, but is imposed for ease of exposition and to maintain symmetry between the assumptions on the different error terms.

above. We record the result here for the empirically relevant case $2K \leq H$. The identification result for the case where $2K > H$ will require further assumptions on the second and higher moments of the error terms (as in Assumption (3)).

Lemma 3 *Suppose that $z_c \perp f_c$, the moments on the right hand side of (3) exist, and Assumption (1) and (4) hold and $2K \leq H$. Then, the parameter vector β is identified.*

3.2 Identification of Specification (D3)

Following the lines of the previous argument (and assuming for simplicity that $K=2$ and $M=1$), under Assumption (1) and if $z_c \perp f_c$, the parameter of interest

$$\beta = (\mathbb{E}(f(z_c) f(z_c)'))^{-1} \mathbb{E}(f(z_c) y_{ch})$$

where

$$f(z) = (\tilde{z}_2, \sin(z), \cos(z))' \tag{5}$$

The only new objects in these moment matrices are $\mathbb{E}(f(z) (\sin(z), \cos(z)))$ and $\mathbb{E}((\sin z, \cos z)' y)$. In a more general setting, Schennach (2004) identifies these moments by identifying the characteristic function of the unobserved variable z and then applying Parseval's Identity. This approach can be adapted to the panel data set context in much the same way as the Hausman, Newey, Ichimura, and Powell (1991) approach: by exploiting the independence of the error terms across households within a cluster.

For expositional ease, assume that there are only two households per cluster. First, since we assume γ is identified, then the unit values $\{v_1, v_2\}$ are also identified and therefore their joint characteristic function ($\mathbb{E}(\exp(i(t_1 v_1 + t_2 v_2)))$) is identified. Substituting into the characteristic function using (2) and using the fact that $\eta_h \perp z$ we obtain

$$\mathbb{E}(\exp(i(t_1 v_1 + t_2 v_2))) = \mathbb{E}(\exp(iz(t_1 + t_2))) \mathbb{E}(\exp(it_1 \eta_{11})) \mathbb{E}(\exp(it_2 \eta_2))$$

Taking logs on both sides we obtain

$$\Pi_{v_1, v_2}(t_1, t_2) = \Pi_z(t_1 + t_2) + \Pi_{\eta_1}(t_1) + \Pi_{\eta_2}(t_2)$$

where $\Pi_x(s)$ denotes the log of the characteristic function of the random variable x evaluated at the point s . Taking the derivative on both sides of the equation above with respect to t_1 and evaluating the derivative function at $t_1 = 0$ and using the fact that the mean of η_2 is zero, we obtain

$$\left. \frac{d}{dt_1} \Pi_{v_1, v_2}(t_1, t_2) \right|_{t_1=0} = \left. \frac{d}{dt_1} \Pi_z(t_1 + t_2) \right|_{t_1=0} \tag{6}$$

the last expression is the derivative of the log of the characteristic function of z evaluated at t_2 . Define the left hand side (which is identified) to be $\phi(t_2)$. We can then use the fundamental

theorem of calculus and obtain the characteristic function of z , $\psi(s)$ as

$$\psi(s) \equiv \exp(\Pi_z(s)) = \exp\left(\int_0^s \phi(t_2) dt_2\right) = \exp\left(\int_0^s \frac{\mathbb{E}[iv_1 \exp(it_2 v_2)]}{\mathbb{E}[\exp(it_2 v_2)]} dt_2\right)$$

assuming enough smoothness to interchange the differentiation and integration operations. Once the characteristic function ψ is known, $\mathbb{E}(q(z))$ for any (measurable) function $q(\cdot)$ can be calculated most cleanly using Parseval's Identity

$$\mathbb{E}(q(z)) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \Xi(q(\cdot), -s) \psi(s) ds \quad (7)$$

where $\Xi(q(\cdot), s) = \int q(t) \exp(its) dt$ is the Generalized Fourier Transform (GFT) of the function q . The GFTs of polynomial and trigonometric functions (and their products) are relatively easy to calculate and are given by the Dirac "Delta" Function (denoted by $\delta(s)$) and its derivatives⁵ evaluated at a finite set of points. To illustrate,

$$\begin{aligned} \mathbb{E}(\sin(z)) &= \frac{1}{2\pi} \int \Xi(\sin(\cdot), -s) \psi(s) ds \\ &= \frac{i}{2\pi} (\psi(1) - \psi(-1)) \end{aligned}$$

A complete list of the relevant GFTs is available upon request. For all the functions $q(\cdot)$ considered here, the integral in (7) reduce to the evaluation of the characteristic function ψ and its derivatives at a finite set of points. Parallel to the argument above, the expectations $\mathbb{E}(y f_j(z))$ for an element $f_j(z)$ of the vector $f(z)$ are similarly identified and the formula is

$$\mathbb{E}(y f_j(z)) = \frac{-i}{2\pi} \int_{-\infty}^{\infty} \Xi(f(\cdot), -s) \psi(s) \frac{\mathbb{E}(i \exp(isv_2) y_1)}{\mathbb{E}(\exp(isv_2))} ds \quad (8)$$

and as in the preceding argument, this calculation simplifies because of the relatively simple form of the GFTs of the functions being considered.

4 Estimation and Large Sample Distribution

The estimation strategy follows the identification argument closely. We first construct the sample versions of the population parameters which for convenience we define as

$$\theta \equiv \left(\{\zeta_j\}_{j=1}^{2K}, \{\xi_j\}_{j=1}^K \right) \quad (9)$$

. We first discuss the estimation of the ζ_j parameters (the discussion for estimation of ξ_j is similar). First, If we knew the value of γ_2 , an unbiased and consistent estimator of $\mathbb{E}(z_c^j)$ would be the

⁵See for instance [Lighthill \(1996\)](#) for a relatively formal treatment of these objects (also see [Schemach \(2004\)](#)).

statistic $\tilde{\zeta}_j$ given by

$$\tilde{\zeta}_j = \frac{1}{C} \sum_{c=1}^C \left[\frac{1}{\binom{H}{j}} \sum_{h_1=1}^{H-j+1} \sum_{h_2=h_1+1}^{H-j+2} \cdots \sum_{h_j=h_{j-1}}^H v_{ch_1} v_{ch_2} \cdots v_{ch_j} \right]$$

However, since in practice γ_2 is estimated (see below for more details), we construct $\hat{v}_{ch} = u_{ch} - x_{ch}' \hat{\gamma}_2$ and the estimator is given by

$$\hat{\zeta}_j = \frac{1}{C} \sum_{c=1}^C \left[\frac{1}{\binom{H}{j}} \sum_{h_1=1}^{H-j+1} \sum_{h_2=h_1+1}^{H-j+2} \cdots \sum_{h_j=h_{j-1}}^H \hat{v}_{ch_1} \hat{v}_{ch_2} \cdots \hat{v}_{ch_j} \right] \quad (10)$$

Similar arguments lead to

$$\hat{\xi}_j = \frac{1}{C} \sum_{c=1}^C \left[\frac{1}{\binom{H}{j}} \sum_{h_1=1}^{H-j+1} \sum_{h_2=h_1+1}^{H-j+2} \cdots \sum_{h_j=h_{j-1}}^H \hat{v}_{ch_1} \hat{v}_{ch_2} \cdots \hat{v}_{ch_{j-1}} \hat{y}_{ch_j} \right] \quad (11)$$

where $\hat{y}_{ch} = \ln q_{ch} - x'_{ch} \hat{\gamma}_2$ for the first specification and $\hat{y}_{ch} = w_{ch} - x'_{ch} \hat{\gamma}_2$ for the second specification. In an additional appendix, we outline the asymptotic properties of the estimator $\hat{\theta} = \left(\left\{ \hat{\zeta}_j \right\}_{j=1}^{2K}, \left\{ \hat{\xi}_j \right\}_{j=1}^K \right)$ and show that under a standard set of conditions (for instance as contained in Section 7 of [Newey and McFadden \(1994\)](#))

$$\sqrt{C} (\hat{\theta} - \theta) \Rightarrow \mathcal{N}(0, V_\theta)$$

For the first two specifications, the object of interest β is a smooth function only of θ ⁶ given by

$$q(\theta) = \begin{bmatrix} \zeta_0 & \zeta_1 & \cdot & \zeta_K \\ \zeta_1 & \cdot & \cdot & \zeta_{K+1} \\ \cdot & \cdot & \cdot & \cdot \\ \zeta_K & \cdot & \cdot & \zeta_{2K} \end{bmatrix}^{-1} \begin{bmatrix} \xi_0 \\ \cdot \\ \cdot \\ \xi_K \end{bmatrix}$$

so that we can estimate β by

$$\hat{\beta} = q(\hat{\theta})$$

An application of the ‘‘Delta’’ method therefore suffices to yield the asymptotic normality of the estimator $\hat{\beta}$ with asymptotic variance given by

$$V_\beta = q'(\theta) V_\theta (q'(\theta))^T$$

For the third specification, in addition to θ we also need to estimate the moments $\mathbb{E}(\tilde{z}_2(\sin z, \cos z))$, $\mathbb{E}((\sin z, \cos z)'$ and $\mathbb{E}((\sin z, \cos z)' y)$. Following the identification argument, we estimate the characteristic func-

⁶where $\zeta_0 = 1$ and $\xi_0 = \mathbb{E}(y_{ch})$

tion of the price variable z_c and then use Parseval's identity to compute the required moments. Recall that the characteristic function was given by $\psi(t) = \exp\left(\int_0^t \phi(s) ds\right)$, where the function ϕ is defined in (6) and is estimated by

$$\hat{\phi}(s) = \frac{\sum_{c=1}^C (\hat{v}_{c1} \exp(is\hat{v}_{c2}))}{\sum_{c=1}^C [\exp(is\hat{v}_{c2})]}$$

where I have assumed for simplicity that each cluster has only 2 households. In practice, we can compute the numerator and the denominator by taking all possible $\binom{H}{2}$ distinct combinations of the households to form $\hat{\phi}$. Finally, the characteristic function

$$\hat{\psi}(t) = \exp\left(\int_0^t \hat{\phi}(s) ds\right)$$

is obtained by solving an ordinary differential equation (with the boundary condition that $\psi(0) = 1$). With $\hat{\psi}$ in hand we can estimate any element $\mathbb{E}(q_{jk}(z))$ of the outer product $\mathbb{E}(f(z)f(z)')$ (recall that $f(z)$ was defined in (5)) as

$$\mathbb{E}[\widehat{q_{jk}(z)}] = \frac{1}{2\pi} \int_{-\infty}^{\infty} \Xi(q_{jk}(\cdot), -s) \hat{\psi}(s) ds \quad (12)$$

Since these functions q_{jk} are much simpler than the kinds considered by Schennach (2004), I can compute $\Xi(q_{jk}(\cdot), -s)$ analytically rather than adopt the numerical integration routine suggested in her paper. As discussed in the identification section, evaluating the integral $\mathbb{E}(q_{jk}(z))$ reduces to evaluating the derivatives of the characteristic function ψ at a finite number of points, so that in the sample we compute the expectations by evaluating the sample counterparts of these derivatives. This results in a considerable simplification of the problem. To illustrate, the object $\mathbb{E}(\sin z)$ can be estimated by $(\hat{\psi}(1) + \hat{\psi}(-1)) (i/2)$.

We can estimate $\mathbb{E}(yf_j(z))$ in a similar fashion and the formula is given by (see Schennach (2004) for more details)

$$\mathbb{E}[\widehat{yf_j(z)}] = \frac{-i}{2\pi} \int_{-\infty}^{\infty} \Xi(f(t), t, -s) \hat{\psi}(s) \frac{\sum_{c=1}^C (i \exp(isv_{c2}) y_{c1})}{\sum_{c=1}^C (\exp(isv_{c2}))} ds \quad (13)$$

Here again, because of the polynomial-trigonometric choice of functions, the Fourier transforms take on particularly simple forms (see Appendix) and the problem reduces in the same manner as above. In fact, all the estimates of elements of the matrices in the normal equations can be expressed as functions of the characteristic function and its derivatives evaluated at a finite set of points.

Given estimates of the moments above, we can form estimates of the matrices $\mathbb{E}(f(z)f(z)')$ and $\mathbb{E}(f(z)y)$ and estimate the parameter of interest by

$$\hat{\beta} = \mathbb{E}(\widehat{f(z)f(z)'})^{-1} \mathbb{E}(\widehat{f(z)y}) \quad (14)$$

where the elements of $\mathbb{E}(\widehat{f(z)} f(z))'$ are given by (12) and of $\mathbb{E}(\widehat{f(z)} y)$ by (13). As mentioned earlier, each element of these matrices is a function of the estimated characteristic function and its derivatives evaluated at a finite set of points, that is to say $\left\{ \frac{\partial \hat{\psi}^k(s_0)}{\partial s^k} \right\}_{k=0}^4$ for $s_0 \in \{-C_{K,M}, -1 \dots 1, C_{K,M}\}$.⁷ Further, it is possible to express these derivatives themselves as functions of $(\hat{\psi}(s_0), \hat{\phi}(s_0), \hat{\phi}'(s_0), \dots, \hat{\phi}'''(s_0))$ for $s_0 \in \{-2, -1 \dots 1, 2\}$. Therefore, in order to characterize the limiting distribution of $\hat{\beta}$ above, it is enough to characterize the limiting distribution of these estimated quantities. The details for the asymptotic theory are relegated to the appendix and the conclusion is

$$\sqrt{C}(\hat{\beta} - \beta) \Rightarrow N(0, V)$$

Although $\hat{\beta}$ converges at the regular rate and has a limiting normal distribution, it is difficult to construct a consistent estimator of the variance matrix V because we do not have an explicit formula for it. One reason is that $\hat{\beta}$ is a complicated multistep estimator that depends upon the preliminary steps in a complicated way. In addition, the limiting distribution of some of the preliminary estimates themselves involves variances that are hard to estimate (these distributions are functionals of gaussian processes). We will therefore use the bootstrap. Although proving that the bootstrap works in this model is beyond the scope of this paper, the smoothness of the estimated parameters in the underlying distribution and the (\sqrt{C}) normality result suggest that the bootstrapped standard errors should be consistent.⁸

5 Monte Carlo Results

The estimators themselves are fairly straightforward. The Monte Carlo illustrations serve to illustrate two points: the first is to offer a set of controlled comparisons between the estimators derived in this paper and the estimator proposed in Deaton (1988) in the linear context and the second is to assess the possible effects of misspecification.

5.1 MC for First Specification

The results of the Monte Carlo simulations for the comparisons between the alternative estimators for specification 1 is given in Tables 1 and 2. Both estimators have comparable Mean Squared Errors (MSE) in the linear case, which we would expect to be the case since both are essentially correcting the pooled OLS estimates for the plim bias terms that are present due to measurement error. As robustness checks, varying the correlation between measurement errors or the signal to noise ratio of the mismeasured prices does not affect either estimator asymmetrically either and they both perform well in these situations.

⁷As the notation suggests, the constant $C_{K,M}$ depends upon K and M . For instance with $K = 2$ and $M = 1$, $C_{K,M} = 2$ while for $K = 1$ and $M = 2$, $C_{K,M} = 4$.

⁸Note that strictly speaking this is not enough. See Abadie and Imbens (2006) or counter-example 1 in Bickel and Freedman (1981) for instances of \sqrt{n} consistent and asymptotically normal estimators for which the bootstrap is invalid (at least without further assumptions).

When these estimators are applied to data where the regression function is actually quadratic in prices, both are inconsistent and there is no discernible pattern to be obtained from varying the correlation or signal to noise ratios on the distribution of the estimators. The distribution of the Monte Carlo estimators is wildly off from the truth in both cases. The correctly specified estimator, which takes into account the quadratic form of the regression function, is well behaved as we would expect from the previous section.

Both estimators stem from the same spirit of correcting OLS estimates for measurement error and one might perhaps be able to extend directly the first method to account for non-linearities in the regression function as well.

5.2 MC for Third Specification

The primary purpose of the MC results for the third specification was to provide a check of the \sqrt{C} convergence result (proved in the appendix) for the parameter coefficients. The true model is given by

$$y = 1 + 2z + 4z^2 + 3\sin(z) + 2\cos(z) + \varepsilon$$

where ε is standard normal and the z are unobserved. Instead we observe

$$v_j = z + \eta_j$$

for $j = 1 \dots H$ where η has the Laplace distribution (and is independent of the ε although that is not required). I also report results for the case where η has the normal distribution. The MSE are presented in Table 3. Encouragingly, the Mean Squared Error (MSE) approximately halves when I double the sample size. However, notice that in some of the cases, the MSE falls by much more than would be justified by a central limit theorem argument.

6 Empirical Results

The theory sketched above is a simplification of the data at hand in several respects. The first is that I do not observe positive quantities (and hence any unit values) for a subset of households, and therefore there is a question of how exactly to include such households in the analysis. In the first specification since I work with a log-log specification, I must perforce drop any households with no recorded quantities. In the second and third specifications I include households purchasing no wheat and set their unit value equal to the average unit value for the village they live in. This is a pragmatic imputation with no compelling justification. Second, for the present analysis, I am ignoring cross price effects, although a more complete analysis should include those as well. Third, the data are sampled using a two-stage stratified design, and I ignore weighting issues and treat the sample as if it were independent and identically distributed across clusters. Fourth, there are households that do not purchase any wheat but consume home grown quantities, and one must

take a position on how to treat such observations. Following the literature (e.g. Deaton (1988)), I have used the NSS imputed harvest price of domestic production and calculated the unit value for such households from this imputation.

I work with a sample of 4440 households from rural Maharashtra, of whom 82% had consumed either market-bought or home-grown wheat in the last 30 days. The 3641 households that consumed some wheat form the sample for my first specification. Of the remaining households, 759 lived in villages where at least one other household had consumed some wheat, these are added to the sample for my second and third specifications. The remaining 40 households lived in villages where no one consumed any wheat and are excluded from the analysis. Average total monthly consumption was Rs. 1134 per household, and on average 3.4% of total household expenditure was spent on wheat for those households that purchased a positive quantity. The unit values show that households paid an average of Rs. 5.79 per kilo for wheat. I have not as yet found an independent source for assessing whether these unit values are in fact close to the prevailing prices in rural Maharashtra during the survey, but the assumption for the paper is that these unit values do provide a measure of price, albeit a noisy one.

Table 4 highlights the spatial and temporal variation in unit values. Each column reports the F-statistic from a regression of the (log of the) unit values on a set of regressors. The regressors in the first column are the 5 NSS region dummies and in the second column dummies for the four quarters (Jul-Sept, Oct-Dec, Jan-Mar, Apr-Jun). The F-statistics are all significant at conventional levels and provide some evidence that there is considerable price variation across space and time in our data. Finally, the last column reports the F-statistic from a regression onto a set of village dummies. Since our identification assumption is one of identifying the demand function through between village variation in price, it is encouraging to see that the dummies together are jointly significant in the regression. Finally, Table 5 presents the coefficients from a regression on region and subround dummies.

Table 6 shows the results from the first stage estimation for the various specifications. For each specification, the first stage estimates the parameters $\{\gamma_1, \gamma_2\}$ using a standard fixed effect model where the $x = \{\text{age-sex variables, log(total expenditure), religion, caste status, household size, labour type}\}$. As might be expected, household size and total expenditure turn out to be important regressors. The pattern of results are consistent with Engel’s law in the sense that although richer households purchase more wheat (the second row of coefficients), the share of wheat in the total budget actually declines as households get richer (the second set of coefficients). Of course, the poorest households do not purchase any wheat but the result is robust to their inclusion as well (the final row of coefficients). This does suggest however that the relationship between budget share of wheat and total expenditure is not necessarily monotonic.

Figure 1 depicts the relationship between purged (log) quantities and unit values using local polynomial kernel regression. The function is not clearly linear, and the object of the first specification is to evaluate the implications of approximating the true regression function (recall that the displayed regression line is not of interest since it is the regression on mismeasured prices) solely

with a polynomial specification.

Following the strategy outlined in the identification section above, I next construct sample moments using the purged unit values and quantities to obtain our estimates for the unobserved moments. For instance to estimate the mean of the unobserved price I take the average of the observed unit values across all households. Higher moments are estimated by taking all possible combinations of households that are consistent for the moment of interest and taking their weighted average (where the weights are given by the relative fraction of observations for a particular sample moment). For instance in calculating $E[z^2]$ we use all $\binom{H}{2}$ possible choice of $i \neq j$ and compute $E_{F_c} v_i v_j$. Since some villages may not have H households, the product $v_{ci} v_{cj}$ may not exist for all clusters, and the average is, naturally, over all non-missing values.

The first specification does not use households that report zero quantities. This leaves me with a total of 3641 households spread across 440 villages. I estimate a series of possible models starting with the linear model. In this case, I can compare my estimates with estimates obtained from the method outlined in Deaton (1997). The results are displayed in Table (7). The point estimates for the elasticity are not too dissimilar in both cases, with the Deaton method slightly larger than the linear specification of our method. The most important difference is in the bootstrapped standard errors which are much larger under the newer method so that the corresponding confidence intervals are also much larger (I report the 2.5th and 97.5th percentile based on 1000 bootstrapped resamples at the cluster level for the second stage of estimation).

For models with higher order polynomials

$$\frac{\partial}{\partial \ln p} \mathbb{E}[q | \ln p] \tag{15}$$

is a function of the price, and one has to take a position on where to evaluate these derivatives. I calculate both the sample version of the average derivative

$$\mathbb{E} \left(\frac{\partial}{\partial \ln p} \mathbb{E}[q | \ln p] \right)$$

and in addition also report results for the derivative evaluated at the 25th and 75th percentile of the purged unit value distribution. I estimate polynomials of order up to five although the magnitude of the bootstrapped standard errors for specifications higher than the quadratic call into question any conclusions that one could derive from the higher order polynomial models. Specifically, in these models, we cannot reject the null that the observed elasticity (evaluated at the candidate values of the price variable) is not significantly different from zero.

The specifications that are non-linear in prices seem to provide better approximations to the regression function graphed above, in that the elasticities are lower at the lower end of the distribution and higher at the higher end of the distribution. Note that the plot above is the regression function for quantities conditional on unit values (mismeasured prices), so its usefulness as a guide for the regression function of quantities on the true prices may be limited.

Figure 2 presents the locally weighted quadratic regression of purged budget shares on purged

unit values. The lines correspond to the 5th and 95th percentiles of the distribution for purged unit values. Again, the relationship looks very similar to the relationship in Figure 1. Specifications 2 and 3 attempt to model the relationship between these purged budget shares and the true (unobserved) prices using a polynomial model and a Fourier flexible form approach.

The results from the polynomial models are presented in Table (8). The elasticity estimated in these models (apart from the linear model) is a function of the budget share and the price. I fix the budget share equal to the average budget share for the sample and evaluate the elasticity for two different values of the price variable (the 25th and the 75th percentile of the purged unit values) as well as the average derivative calculation. As might be suggested by Figure 2, the elasticities are lower at the 25th percentile of the purged unit values and higher when evaluated at the 75th percentile of the unit value distribution. The average elasticity (evaluated at the mean of the purged unit value distribution) lies between these two. All the models in Table 8 exhibit this pattern. However, note that the standard errors become extremely large after the cubic specification, and indeed for all but one of the quartic and quintic specifications we cannot reject the null of zero elasticity. The large standard errors for the higher degree specifications reflect the imprecision with which the higher moments are estimated (as reflected in their bootstrapped standard errors which are not reported here). Finally, the results from the last specification (setting $K=2$ and $M=2$) are displayed in Table 9. The point estimates are broadly similar to the ones from the previous specification and here again the elasticities are lower at the lower end of the price distribution relative to the higher end.

Looking across all the specifications we see that the results across various non-linear models are broadly consistent with each other. While there is no compelling reason to believe any of the regression functions is correctly specified, the relative stability of the elasticity estimates across specifications is encouraging and suggests that misspecification is not driving the results. Overall, the elasticities derived from the budget share specification are more precisely estimated (as expected) relative to the log-quantity specification. While no specification clearly dominates the others, the quadratic choice for all three specifications is relatively straightforward to implement and provides broadly consistent and reasonably precise estimates of the price elasticities.

For comparison, we provide results from estimating a specification without accounting for measurement error. We implement a quadratic version of the first specification and the results are presented in Table (10) and should be compared against the third row of results in Table (7). Both the point estimates and the standard errors for the naive estimator differ from those of the estimator that accounts for measurement error. Two differences emerge: the estimates without correction suggest much less sampling uncertainty than the results in Table (7) which is reasonable since the former do not account for measurement error in any way. If the proposed model in the paper is correct, the naive estimator seems to significantly underestimate sampling uncertainty. Second, the point estimates are quite different as well and suggest much smaller differences in the elasticities across the price distribution. The differences in this case seem to stem from the fact that the estimated higher moments of the purged price distribution are quite different for the two

models. If specification (1) is correct, then the naive estimator's computation of these moments will be inconsistent since it does not purge the measurement error.

To conclude, the most substantive point and one that comes across in all the specifications, is that there is evidence that the price elasticity does vary across the price distribution. Estimated price elasticities are generally lower (in absolute magnitude) in the lower tails of the distribution and are generally larger in the upper tails, something that a linear specification of the demand equation would miss.

7 Conclusions

This paper attempts to estimate price elasticities by modeling flexibly the regression function for quantities demanded (and budget shares) as a function of price while at the same time accounting for the effect of measurement error in prices that is fundamentally non-classical in nature. In such a setting, standard instrumental variable techniques are no longer available. We adapt the strategy suggested by [Hausman, Newey, Ichimura, and Powell \(1991\)](#) and [Schennach \(2004\)](#) to account for the non-classical nature of the measurement error and exploit the clustered nature of the data to achieve identification of the price elasticity. We next propose an estimator based on the identification strategy and discuss its large sample properties. Next, we discuss its performance in Monte Carlo simulations and finally apply it to estimate a price elasticity using data from rural India. Kernel regressions from the data suggest potential evidence of non-linearity in the true conditional expectation and we estimate a series of alternative specifications to account for this. The estimator confirms the initial presumption and for all the specifications considered the estimated price elasticities are somewhat lower at the bottom of the price distribution and higher at the top.

Two potential further directions for research are to include and estimate cross-price effects and also to consider estimation in the case where the number of polynomial (or trigonometric) terms increases with the sample size.

A Appendix

A.1 Identification

Proof of Lemma 1.

Proof. Consider the directly identified quantity (for $h_a \neq h_b$ for all $(a, b) \in \{1, \dots, j\}$)

$$\begin{aligned} \mathbb{E}(v_{ch_1} \dots v_{ch_j}) &= \mathbb{E}\left((z_c + \eta_{ch_1}) \dots (z_c + \eta_{ch_j})\right) \\ &= \mathbb{E}\left(z_c^j + z_c^{j-1} \sum_{h_1=1}^j \eta_{ch_1} + \dots + z_c \sum_{h_1=1}^2 \sum_{h_{j-1}=h_{j-2}+1}^j \left(\eta_{ch_1} \dots \eta_{ch_{j-1}}\right) + \left(\eta_{ch_1} \dots \eta_{ch_j}\right)\right) \\ &= \mathbb{E}(z_c^j) \end{aligned}$$

where the last equality follows from the independence between z_c and η_{ch} and the fact that $\mathbb{E}(\eta_{ch}) = 0$

(which are Assumption (2)). Note that for this result alone, mean independence rather than independence will suffice. Further, we can consider all $\binom{H}{j}$ distinct subsets of size j from the set $\{v_1, \dots, v_H\}$ each of whose expectation will be equal to that of $\mathbb{E}(z_c^j)$ by the same argument as above so that

$$\mathbb{E}(z_c^j) = \mathbb{E} \left[\frac{1}{\binom{H}{j}} \sum_{h_1=1}^{H-j+1} \sum_{h_2=h_1+1}^{H-j+2} \dots \sum_{h_j=h_{j-1}}^H v_{ch_1} v_{ch_2} \dots v_{ch_j} \right]$$

■

Proof of Lemma 2

Proof. As before in Lemma 1, consider $j+1$ distinct households and the directly identified quantity $\mathbb{E}(v_{ch_1} \dots v_{ch_j} y_{ch_{j+1}})$. Rewriting this we obtain

$$\begin{aligned} & \mathbb{E} \left(y_{ch_{j+1}} \left(z_c^j + z_c^{j-1} \sum_{h_1=1}^j \eta_{ch_1} + \dots + z_c \sum_{h_1=1}^2 \sum_{h_{j-1}=h_{j-2}+1}^j \left(\eta_{ch_1} \dots \eta_{ch_{j-1}} \right) + \left(\eta_{ch_1} \dots \eta_{ch_j} \right) \right) \right) \\ &= \mathbb{E}(z_c^j y_{ch_{j+1}}) + \mathbb{E} \left(y_{ch_j} z_c^{j-1} \sum_{h_1=1}^j \eta_{ch_1} \right) + \mathbb{E} \left(y_{ch_j} \sum_{h_1=1}^2 \sum_{h_{j-1}=h_{j-2}+1}^j \left(\eta_{ch_1} \dots \eta_{ch_{j-1}} \right) \right) \\ & \quad + \mathbb{E} \left(y_{ch_j} \left(\eta_{ch_1} \dots \eta_{ch_j} \right) \right) \\ &= \mathbb{E}(z_c^j y_{ch_{j+1}}) \end{aligned}$$

where the last equality follows from the fact that all the terms are of the form

$$\mathbb{E} \left(y_{ch_{j+1}} g(z_c) r \left(\eta_{ch_1}, \dots, \eta_{ch_j} \right) \right)$$

for appropriately specified functions $g(\cdot)$ and $r(\cdot)$. This expectation is identically zero since

$$\begin{aligned} & \mathbb{E} \left(y_{ch_{j+1}} g(z_c) r \left(\eta_{ch_1}, \dots, \eta_{vh_j} \right) \right) = \\ & \mathbb{E} \left((f(z_c) + f_c + \varepsilon_{ch_{j+1}}) g(z_c) r \left(\eta_{ch_1}, \dots, \eta_{vh_j} \right) \right) \\ &= \mathbb{E} \left(f(z_c) g(z_c) r \left(\eta_{ch_1}, \dots, \eta_{vh_j} \right) \right) + \mathbb{E} \left(f_c g(z_c) r \left(\eta_{ch_1}, \dots, \eta_{ch_j} \right) \right) + \mathbb{E} \left(\varepsilon_{ch_{j+1}} r \left(\eta_{ch_1}, \dots, \eta_{ch_j} \right) \right) \\ &= 0 \end{aligned}$$

where the first two terms are zero since $(\eta_{ch_s})_{s=1}^j$ is independent of (z_c, f_c) and $\mathbb{E} \left(r \left(\eta_{ch_1}, \dots, \eta_{ch_j} \right) \right) = 0$ and the last term is zero since $\varepsilon_{ch_{j+1}}$ is independent of $(\eta_{ch_s})_{s=1}^j$ (all of which follow from Assumption 4) ■

Proof of Lemma 3

Proof. Under Assumption (1), the parameters $\gamma = (\gamma_1, \gamma_2)$ are identified and therefore $\{y_{ch}, v_{ch}\}$ are identified. Next, under the assumptions stated in the lemma β can be obtained as

$$\beta = \mathbb{E}(\tilde{z}_{cK} \tilde{z}'_{cK})^{-1} \mathbb{E} \left(\frac{1}{H} \sum_{h=1}^H \tilde{z}_{cK} y_{ch} \right)$$

and each element on the right hand side is identified by the previous two lemmas. ■

A.2 Large Sample Theory

The asymptotic theory for the first two specifications is straight-forward and can be derived using standard results from two-step estimation theory (for instance Section 6 of [Newey and McFadden \(1994\)](#)). In the first step, the elements of γ can be consistently estimated (upto a constant) using the sample version of the moment conditions⁹

$$\mathbb{E} \left(\underbrace{\begin{pmatrix} \sum_{h=1}^H (\ddot{w}_{ch} - \ddot{x}'_{ch} \gamma_1) \ddot{x}_{ch} \\ \sum_{h=1}^H (\ddot{u}_{ch} - \ddot{x}'_{ch} \gamma_1) \ddot{x}_{ch} \end{pmatrix}}_{m((w_{ch}, x_{ch}, u_{ch})_{h=1}^H, \gamma)} \right) = 0$$

Under the assumptions for Lemma 3 and that the second moments of the bivariate vector of the composite error terms $(\ddot{\varepsilon}_{ch}, \ddot{\eta}_{ch})$ exist¹⁰, standard arguments yield

$$\sqrt{C}(\hat{\gamma} - \gamma) \Rightarrow \mathcal{N}(0, V_\gamma)$$

where

$$V_\gamma = \left(\mathbb{I}_2 \otimes \sum_{h=1}^H \mathbb{E}(\ddot{x}_h \ddot{x}'_h) \right)^{-1} M \left(\mathbb{I}_2 \otimes \sum_{h=1}^H \mathbb{E}(\ddot{x}_h \ddot{x}'_h) \right)^{-1}$$

where M is the second moment matrix of the moment conditions.

In the second step, we estimate the elements of the parameter θ (defined in (9)) by (10) and (11). Rewriting these to reflect the dependence upon the first stage estimates γ and writing them out in a common format

$$\begin{aligned} \frac{1}{C} \sum_{c=1}^C \left[\frac{1}{\binom{H}{j}} \sum_{h_1=1}^{H-j+1} \sum_{h_2=h_1+1}^{H-j+2} \dots \sum_{h_j=h_{j-1}}^H (u_{ch_1} - x'_{ch_1} \hat{\gamma}_2) \dots (u_{ch_j} - x'_{ch_j} \hat{\gamma}_2) \right] - \hat{\zeta}_j &= 0 \\ \frac{1}{C} \sum_{c=1}^C \left[\frac{1}{\binom{H}{j}} \sum_{h_1=1}^{H-j+1} \sum_{h_2=h_1+1}^{H-j+2} \dots \sum_{h_j=h_{j-1}}^H (u_{ch_1} - x'_{ch_1} \hat{\gamma}_2) \dots (u_{ch_{j-1}} - x'_{ch_{j-1}} \hat{\gamma}_2) (w_{ch_j} - x'_{ch_j} \hat{\gamma}_1) \right] - \hat{\xi}_j &= 0 \end{aligned}$$

where for the first specification we replace w_{ch} with q_{ch} . We collect these $2K + K$ moment conditions and denoting them collectively by the moment vector $g((w_{ch}, x_{ch}, u_{ch})_{h=1}^H, \hat{\theta}, \hat{\gamma})$. Note that the moment conditions are in fact linear in the second stage parameters. Under the assumptions above and the additional assumption

ASSUMPTION 5 *The error terms $(\ddot{\varepsilon}_{ch}, \ddot{\eta}_{ch})$ have finite second moments. The true parameter (θ, γ) lies in the interior of the euclidean parameter space. The moment function g satisfies $\mathbb{E} \left(\left\| g((w_{ch}, x_{ch}, u_{ch})_{h=1}^H, \theta, \gamma) \right\|^2 \right)$ is finite and its first derivative (with respect to (θ, γ)) is uniformly bounded over a neighbourhood of the truth in expectation. The expressions in the first three displays on the next page exist and the matrix V_θ is non-singular.*

We can now verify the conditions of Theorem 6.1 of [Newey and McFadden \(1994\)](#) to conclude that

$$\sqrt{C}(\hat{\theta} - \theta) \Rightarrow \mathcal{N}(0, V_\theta)$$

⁹Note that no claim of optimality for the choice of moment conditions is being made. It is possible to construct more efficient estimators of γ given the assumptions above, however this is not pursued here. The double dots denote that the variables are in deviations from their cluster means.

¹⁰As usual, the notation $f(z_c)$ denotes the flexible specifications of the price variable in the demand equation.

where

$$V_\theta = \mathbb{E}(rr')$$

where

$$r = g\left((w_{ch}, x_{ch}, u_{ch})_{h=1}^H, \theta, \gamma\right) + G_\gamma m\left((w_{ch}, x_{ch}, u_{ch})_{h=1}^H, \gamma\right)$$

and

$$G_\gamma = \mathbb{E}\left(\frac{\partial g\left((w_{ch}, x_{ch}, u_{ch})_{h=1}^H, \theta, \gamma\right)}{\partial \gamma}\right)$$

A.3 Asymptotic Distribution for the Third Specification

Under the assumptions outlined above, we derive the limiting distribution for the case where $K=2$ and $M=2$. As mentioned in the text, in order to characterize the limiting distribution of the estimator $\hat{\beta}$, it suffices to characterize the limiting distribution of the vector

$$\left(\hat{\psi}(s_0), \hat{\phi}(s_0), \hat{\phi}'(s_0), \dots, \hat{\phi}'''(s_0)\right) \quad (16)$$

for $s_0 \in \{-4, -1, \dots, 1, 4\}$. Apart from the first term, each term can be estimated by its sample average; this is because we can calculate the derivatives of $\frac{\mathbb{E}[v_1 \exp(itv_2)]}{\mathbb{E}[\exp(itv_2)]}$ analytically, invoking the appropriate DCT to allow interchange of integration and differentiation. Each derivative can then be consistently estimated by the analogy principle and the analysis of the limiting distribution is considerably simplified by the fact that both the numerator and denominator are Lipschitz continuous in t . However, there is an added complication due to the fact that we do not observe v_s directly. Consider for instance the estimator of the parameter

$$\phi(t) = \frac{\mathbb{E}[v_1 \exp(itv_2)]}{\mathbb{E}[\exp(itv_2)]}$$

which we estimate by

$$\hat{\phi}(t) = \frac{\sum_{c=1}^C (\hat{v}_{c1} \exp(it\hat{v}_{c2}))}{\sum_{c=1}^C \exp(it\hat{v}_{c2})} \equiv \frac{\hat{m}_1(t)}{\hat{m}_2(t)}$$

In order to study the asymptotic properties of this estimator, we first introduce an easier (but infeasible) estimator

$$\tilde{\phi}(t) = \frac{\frac{1}{C} \sum_{c=1}^C v_{c1} \exp(itv_{c2})}{\frac{1}{C} \sum_{c=1}^C \exp(itv_{c2})} \equiv \frac{\tilde{m}_1(t)}{\tilde{m}_2(t)}$$

We will study the limiting properties of $\sqrt{C}(\hat{\phi}(t) - \phi(t))$ by looking at

$$\sqrt{C}(\hat{\phi}(t) - \phi(t)) = \sqrt{C}(\hat{\phi}(t) - \tilde{\phi}(t)) + \sqrt{C}(\tilde{\phi}(t) - \phi(t))$$

and we will characterize the joint limiting properties of $\sqrt{C}(\hat{\phi}(t) - \tilde{\phi}(t))$ and $\sqrt{C}(\tilde{\phi}(t) - \phi(t))$.

Consider first the second term $\sqrt{C}(\tilde{\phi}(t) - \phi(t))$ and consider first the convergence properties of the numerator $\tilde{m}_1(t)$ and the denominator $\tilde{m}_2(t)$ individually (we will then apply the delta method to obtain the limiting properties of $\tilde{\phi}$). Standard arguments can be used to show that for a fixed t , the normalized sample averages $\sqrt{C}(\tilde{m}_1(t) - m_1(t), \tilde{m}_2(t) - m_2(t))' \equiv (\sqrt{C}(\tilde{m}(t) - m(t)))$ converge in distribution to a bivariate normal distribution. In fact, they can be used to show that for any finite set of $\{t_1, \dots, t_k\}$ the corresponding

vector $\sqrt{C}((\tilde{m}(t_1) \dots \tilde{m}(t_k))' - (m(t_1) \dots m(t_k)))$ converges to a multivariate normal distribution.¹¹ For any fixed t then, the asymptotic normality of $\sqrt{C}(\hat{\phi}(t) - \phi(t))$ then follows from an application of the “Delta” method and the additional assumption that $m_2(t) \neq 0$.

Consider the first term now. We can rewrite the difference between $\hat{m}_2(t)$ and $\tilde{m}_2(t)$ as

$$\frac{1}{\sqrt{C}} \sum_{c=1}^C \exp(it(u_{ch_2} - x'_{ch}\hat{\gamma}_2)) - \exp(it(u_{ch_2} - x'_{ch}\gamma_2))$$

and a standard mean-value argument and the results for the first-stage estimation in (14) will yield that $\sqrt{C}(\hat{m}_2(t) - \tilde{m}_2(t))$ will converge weakly to a normal distribution with zero mean and variance

$$\mathbb{E}(x_{ch} \exp(itv_{ch})) V_{\gamma} \mathbb{E}(x'_{ch} \exp(itv_{ch}))$$

Similar arguments yield the asymptotic normality for $\sqrt{C}(\hat{m}_1(t) - \tilde{m}_1(t))$. The “Delta” method will then yield the asymptotic normality of $\sqrt{C}(\hat{\phi}(t) - \tilde{\phi}(t))$ and hence the asymptotic normality of $\sqrt{C}(\hat{\phi}(t) - \phi(t))$ (note that the variance of the limiting distribution will require an additional calculation of the covariance between the two limit distributions). Similar arguments can be adduced to show the convergence of the expressions $(\hat{\phi}'(s_0), \dots, \hat{\phi}'''(s_0))$.

A.3.1 Large Sample Properties for $\hat{\pi}(s)$

The large sample distribution for the estimator $\hat{\pi}(s)$ is somewhat more complicated since it depends upon the value of the estimator $\hat{\phi}$ at a all points $[0, s]$ rather than just a finite number of them. This in turn implies that we need to consider not just the pointwise convergence of $\hat{\phi}$ (in t) but rather convergence that is uniform in t in some appropriate sense.

As a first step, we will need to show that $\sqrt{C}(\hat{m}(\cdot) - m(\cdot))$ when viewed as stochastic progresses (in t for $t \in [0, s]$) converge weakly. As before, we consider the decomposition

$$\sqrt{C}(\hat{m}(t) - m(t)) = \sqrt{C}(\hat{m}(t) - \tilde{m}(t)) + \sqrt{C}(\tilde{m}(t) - m(t)) \quad (17)$$

We first use results from the empirical process literature to study the limiting properties of the second term. One method is to show that the function classes (suppressing the dependence upon h)

$$\mathcal{F} = \{t \in [0, s] : f(v_c, t) = \exp\{iv_c t\}\}$$

$$\mathcal{G} = \{t \in [0, s] : f(v_{1c}, v_{2c}, t) = v_{1c} \exp\{iv_{2c} t\}\}$$

are Donsker. In particular, it is easy to show that conditions of Example 19.7 of [van der Vaart \(1998\)](#) are satisfied since the functions are Lipschitz in t with appropriate moments existing because of the identification assumptions. Therefore we can conclude that the class \mathcal{F} above is Donsker and hence converge weakly to a tight (Gaussian) process W_1 with zero mean covariance given by

$$\text{Cov}(W_1(s), W_1(t)) = \mathbb{E}(\exp(i(t+s)v_c)) - \mathbb{E}(\exp(itv_c)) \mathbb{E}(\exp(isv_c))$$

(see also [Feuerverger and Mureika \(1977\)](#) for more details). Similarly, we can show that the class \mathcal{G} is also

¹¹In fact, we will show below that the process (indexed by $t \in [0, s]$, $\sqrt{C}(\tilde{m}(\cdot) - m(\cdot))$) converges to a Gaussian process because that is needed to study the limiting distribution of $\hat{\psi}$.

Donsker and also converges weakly to a tight Gaussian process W_2 with zero mean and covariance given by

$$Cov(W_2(s), W_2(t)) = \mathbb{E}(v_{1c}^2 \exp(i(t+s)v_{2c})) - \mathbb{E}(v_{1c} \exp(itv_{2c})) \mathbb{E}(v_{1c} \exp(isv_{2c}))$$

Since showing the Donsker property element by element implies the Donsker property for the vector of elements, we can conclude that the process $\sqrt{C}(\tilde{m}(\cdot) - m(\cdot))$ converges weakly to to Gaussian limit process $X(\cdot)$.

Finally, we deal with the first term in the decomposition (17). Consider the term

$$\begin{aligned} \tilde{W}_{1C}(t) &= \hat{m}_2(t) - \tilde{m}_2(t) \\ &= \frac{1}{\sqrt{C}} \sum_{c=1}^C \exp(it(u_{ch} - x'_{ch}\hat{\gamma}_2)) - \exp(it(u_{ch} - x'_{ch}\gamma_2)) \end{aligned}$$

Using the previous results it follows that for any finite collection $(t_s)_s$ the vector $(\tilde{W}_{1C}(t_s))_s$ converges to a normal distribution \tilde{W}_1 with zero mean and covariance given by

$$Cov(\tilde{W}_1(t_a), \tilde{W}_1(t_b)) = \mathbb{E}(\exp(it_a v_{ch}) x'_{ch}) V_\gamma \mathbb{E}(\exp(it_b v_{ch}) x_{ch})$$

In addition, the space $[0, s]$ with the usual metric is totally bounded. In order to show a weak convergence result for the process $\tilde{W}_{1C}(\cdot)$ we therefore only need to verify that condition (ii) (asymptotic tightness) of Theorem 18.14 of [van der Vaart \(1998\)](#) holds. To see this, for any given (ε, η) , partition $[0, s]$ into intervals of length no larger than $\frac{\varepsilon}{c\Phi^{-1}(1-\eta)}$ where Φ^{-1} is the inverse of the standard normal CDF. Then, as $C \rightarrow \infty$ the probability that the maximum value of the difference $\tilde{W}_{1C}(s) - \tilde{W}_{1C}(t)$ over any partition (i.e. (s, t) are in the same partition) will exceed ε will be less than η . Therefore the process is also asymptotically tight and the weak convergence result follows. Similar arguments can be used to show that the process $\sqrt{C}(\hat{m}_1(\cdot) - \tilde{m}_1(\cdot))$ also converges to a Gaussian process. Therefore, by the Continuous Mapping Theorem, each element of the vector valued process $\sqrt{C}(\hat{m}(\cdot) - m(\cdot))$ converges to a Gaussian Process, which we denote by $X(\cdot)$.

Having shown that the process $\sqrt{C}(\hat{m}(\cdot) - m(\cdot))$ converges to a Gaussian process we next apply the Functional ‘‘Delta’’ method to the function

$$g(x(\cdot)) = \int_0^s \frac{x_1(t)}{x_2(t)} dt$$

This function is Frechet Differentiable at $m(\cdot)$ with continuous derivative $\mathcal{L}_m(\cdot)$ given by

$$\mathcal{L}_m(h) = \int_0^s \frac{1}{m_2(t)} h_1(t) - \frac{m_1(t)}{(m_2(t))^2} h_2(t) dt$$

Then, we can apply Theorem 20.8 of [van der Vaart \(1998\)](#) to conclude that $\sqrt{C}(\hat{\pi}(s) - \pi(s))$ converges to the random variable $\mathcal{L}_m(X(\cdot))$ and since $\mathcal{L}_m(\cdot)$ is a continuous linear functional applied to a tight Gaussian process, $\mathcal{L}_m(X(\cdot))$ will be normally distributed (see for instance Lemma 3.9.8 of [van der Vaart and Wellner \(1996\)](#)).

References

- ABADIE, A., AND G. W. IMBENS (2006): “On the Failure of the Bootstrap for Matching Estimators,” National Bureau of Economic Research Cambridge, Mass., USA.
- BICKEL, P. J., AND D. FREEDMAN (1981): “Some Asymptotic Theory for the Bootstrap,” *The Annals of Statistics*, 9(6), 1196–1217.
- DEATON, A. (1987): “Estimation of Own and Cross Price Elasticities from Household Survey Data,” *Journal of Econometrics*, 36, 7–30.
- (1988): “Quality, Quantity and the Spatial Variation of Price,” *American Economic Review*, pp. 418–430.
- DEATON, A. (1997): *The Analysis of Household Surveys: A Microeconomic Approach to Development Policy*. The Johns Hopkins University Press.
- DEATON, A., AND J. MUELLBAUER (1980): “An Almost Ideal Demand System,” *American Economic Review*, 70(3), 312–326.
- DEATON, A., AND S. NG (1998): “Parametric and Non-Parametric Approaches to Price and Tax Policy Reform,” *Journal of the American Statistical Association*, 93, pp.900–909.
- EUBANK, R., AND P. SPECKMAN (1990): “Curve Fitting by Polynomial-Trigonometric Regression,” *Biometrika*, 77, 1–9.
- FEUERVERGER, A., AND R. MUREIKA (1977): “The Empirical Characteristic Function and its Applications,” *Annals of Statistics*, 5, 88–97.
- GALLANT, A. (1981): “On the Bias in Flexible Functional Forms and an Essentially Unbiased Form,” *Journal of Econometrics*, 15, 211–245.
- HARDLE, W., AND T. STOKER (1989): “Investigating Smooth Multiple Regression by the Method of Average Derivatives,” *Journal of the American Statistical Association*, 84(No. 408), pp. 986–995.
- HAUSMAN, J., W. NEWEY, H. ICHIMURA, AND J. POWELL (1991): “Identification and Estimation of Polynomial Errors-in-Variables Models,” *Journal of Econometrics*, 50(3), 273–296.
- KENNAN, J. (1989): “Simultaneous Equations Bias in Disaggregated Econometric Models,” *The Review of Economic Studies*, 56, 151–156.
- LIGHTHILL, M. (1996): *Introduction to Fourier Analysis and Generalized Functions*. Cambridge University Press.
- NEWEY, W., AND D. MCFADDEN (1994): “Large Sample Estimation and Hypothesis Testing,” in *Handbook of Econometrics*, ed. by R. Engle, and D. McFadden, vol. IV, chap. 36, pp. 2111–2245. Elsevier Science.
- SCHENNACH, S. (2004): “Estimation of Non Linear Models with Measurement Error,” *Econometrica*, 72(1), 33–75.
- VAN DER VAART, A. (1998): *Asymptotic Statistics*. Cambridge University Press.
- VAN DER VAART, A., AND J. WELLNER (1996): *Weak Convergence and Empirical Processes With Applications to Statistics*. Springer.

Table 1: Monte Carlo Results (500 simulations, Clusters=300, HH=3)

Error Correlation	-.25		-.5		-.75	
	Deaton	HNIP	Deaton	HNIP	Deaton	HNIP
Method						
Mean ($\beta = 5$)	4.976587	5.003137	4.977571	5.004216	4.974316	5.000928
S.D	.04987	.05058	.05777	.05854	.0524	.052953

Table 2: Monte Carlo Results (500 Simulations,C=300)

Signal Noise	.1		.5		.9	
	Deaton	HNIP	Deaton	HNIP	Deaton	HNIP
Estimator						
Mean ($\beta = 5$)	4.974557	5.000918	4.974139	5.000687	4.975247	5.002013
S.D.	.054903	.0555145	.051808	.0525022	.0533261	.0537687

Table 3: Mean Square Errors for the Monte Carlo Simulations (300 simulations)

Laplace Errors			Normal Errors		
N=1000	N=2000	MSE ₂₀₀₀ /MSE ₁₀₀₀	N=1000	N=2000	MSE ₂₀₀₀ /MSE ₁₀₀₀
292.67	117.7	.402	131.58	55.445	.42138
127.26	12.137	.095	43.515	17.395	.39975
35.303	14.108	.399	19.6973	6.9773	.3584
270.52	25.638	.094	92.082	36.084	.39186
334.46	137.58	.411	141.68	63.561	.44863

Table 4: F- Statistics from a regression of the log of unit values

RHS Variables	Region	Subround	Region*Subround	Village Dummies
F-Statistics	49.83	23.65	18.96	8.6

Table 5: Regional and Seasonal Differences in Unit Values

depvar	R2	R3	R4	R4	R5	SR2	SR3	SR4
ln(unit value)	-.017	-.19	-.159	-.15	-.09	.031	.087	.097
(S.E)	(.015)	(.018)	(.016)	(.016)	(.02)	(.012)	(.013)	(.013)

Table 6: Results from First Stage Regressions. Regressors include log(total expenditure), log(household size) and sociodemographic characteristics.

DepVar	ln(expend)	ln(hhsize)	Hindu	Schedule Caste
Ln(unit value)	.073 (.008)	-.044 (.01)	-.010 (.04)	-.003 (.014)
Ln(quantity) (n=3641)	.354 (.04)	.257 (.05)	-.03 (.17)	.025 (.067)
Budges Share (n=4400)	-.013 (.001)	.012(.002)	.0001 (.008)	-.006 (.002)

Table 7: Average Price Elasticities for different models under specification 1

Model	25 th Percentile	Average Derivative	75 th Percentile
Deaton	-	-1.438 (-1.84,-1.35)	-
Linear	-1.575 (-2.13,-1.22)	-1.575 (-2.13, -1.22)	-1.575 (-2.13,-1.22)
Quadratic	-.5607 (-1.30,.32)	-1.628 (-2.21, -1.26)	-2.393 (-3.59, -1.84)
Cubic	-.6626 (-7.82, 3.52)	-1.723 (-4.24, .52)	-2.536 (-10.47,4.86)
Quartic	-.7785 (-6.86, 5.56)	-1.769 (-5.01, .45)	-2.555 (-13.01, 6.56)
Quintic	-4848 (-7.39, 4.20)	-1.531 (-3.16, -.23)	-2.141 (-8.71, 4.03)

All Confidence Intervals based on 1000 bootstrapped replications

Table 8: Elasticities estimated from Specification 2

Model	25 th Percentile	Average Derivative	75 th Percentile
Deaton	-	-1.465(-2.15, -1.13)	-
Linear	-1.463 (-1.95,-.89)	-1.463 (-1.95,-.88)	-1.463 (-1.95,-.89)
Quadratic	-.965 (-1.58,-.25)	-1.424 (-1.89,-.89)	-1.765 (-2.35, -1.21)
Cubic	-.988 (-1.84,.175)	-1.445 (-2.20,-.69)	-1.803 (-3.30, -.85)
Quartic	11.94 (-13.19,6.74)	-.089 (-2.43,.68)	-10.10 (-6.32, 4.66)
Quintic	-1.28 (-9.31,8.78)	-1.50 (-2.44,-.48)	-1.532 (-6.48, 4.65)

Elasticities are evaluated at the average budget share for three different points in the unit value distribution. All Confidence Intervals based on 1000 bootstrapped replications

Table 9: Elasticities estimated from Specification 3 (evaluated at the average budget share and at three different price values)

25 th Percentile	Average Derivative	75 th Percentile
-1.3724 (-2.09, -.43)	-.8806 (-1.41,-.28)	-1.5943 (-3.40,-.88)

Elasticities are evaluated at the average budget share for three different points in the unit value distribution. All Confidence Intervals based on 200 bootstrapped replications

Table 10: Elasticities estimated without accounting for measurement error

25 th Percentile	Average Derivative	75 th Percentile
-1.4732 (-1.48, -1.46)	-1.7614 (-1.768, -1.75)	-1.9492 (-1.95,-1.942)

Elasticities are evaluated at the average budget share for three different points in the unit value distribution




Figure 1: Locally quadratic regression of purged quantities on purged unit values using a Gaussian kernel with a bandwidth of .2. The vertical lines represent the 5th and 95th percentiles respectively




Figure 2: Locally quadratic regression of purged budget share on purged unit values using a Gaussian kernel with a bandwidth of .2. The vertical lines represent the 5th and 95th percentiles respectively